

**The Centers for Disease Control and Prevention**

**Antibiotic Treatment Dosing Guidelines for  
The National Pharmaceutical Stockpile  
Components**

## Inhalational Anthrax Treatment Protocol

	Initial Therapy	Duration
<b>Adults:</b>  <b>Intravenous</b>	Ciprofloxacin 400 mg BID <sup>1</sup> <b>OR</b>	<b>Estimated for 7 days.</b> (Switch to <u>oral</u> antibiotic therapy when clinically appropriate to complete <b>60-day</b> <sup>2</sup> regimen.)
	Doxycycline 100 mg BID <b>OR</b>	
	Erythromycin 15-20 mg/kg/day in divided doses	
	Penicillin G 20 MU/day in divided doses	
<b>Oral</b>	Ciprofloxacin 500 mg BID <b>OR</b>	<b>Estimated for 60 days</b> <sup>2</sup>
	Doxycycline 100 mg BID	
<b>Children</b> <sup>3</sup>  <b>Intravenous</b>	Ciprofloxacin 15 mg/kg IV Q12hrs <sup>1,4</sup> <b>OR</b>	<b>Estimated for 7 days.</b> (Switch to <u>oral</u> antibiotic therapy when clinically appropriate to complete <b>60-day</b> <sup>2</sup> regimen.)
	Doxycycline <sup>5</sup> : <b>&gt; 8 yrs and &gt; 45 kg:</b> 100 mg BID <b>&gt; 8 yrs and ≤ 45 kg:</b> 2.2 mg/kg/day in 2 divided doses <b>≤ 8 yrs:</b> (same as <b>&gt; 8 yrs and ≤ 45 kg</b> ) <b>OR</b>	
	Erythromycin 15-20 mg/kg/day IV in divided doses	
	Penicillin G 400,000 Units/kg/day in divided doses	
<b>Oral</b>	Ciprofloxacin 15-20 mg/kg Q12 hrs <sup>4</sup> <b>OR</b>	<b>Estimated for 60 days</b> <sup>2</sup>
	Doxycycline <sup>5</sup> : <b>&gt; 8 yrs and &gt; 45 kg:</b> 100 mg BID <b>&gt; 8 yrs and ≤ 45 kg:</b> 2.2 mg/kg BID <b>≤ 8 yrs:</b> same as above	
<b>Pregnancy</b> <sup>3,6</sup>	Same as for non-pregnant adults (the high mortality rate from the infection outweighs the risk posed by the antibiotic)	
	<b>Doxycycline oral not recommended for more than 14 days of therapy.</b>	
<b>Immuno-compromised</b>	Same as for non-immunocompromised adults and children	

1. Therapy with ciprofloxacin may be initiated either as intravenous or oral dosage form. The pharmacokinetics are such that oral ciprofloxacin is rapidly and well absorbed from the GI tract with no substantial loss by first-pass metabolism. Maximum serum concentrations are attained 1-2 hours after oral dosing.
2. Because the potential persistence of spores following a possible aerosol exposure, antibiotic therapy should be continued for at least 30 days if used alone, and although supporting data are less definitive, longer antibiotic therapy (up to 42-60 days) might be indicated.
3. If susceptibility testing allows, therapy should be changed to IV penicillin for treatment or oral amoxicillin for post-exposure prophylaxis to continue therapy out 60 days.
4. Ciprofloxacin dose should not exceed 1gram/day in children.
5. In 1991, the American Academy of Pediatrics amended their recommendation to allow treatment of young children with tetracyclines for serious infections, such as, Rocky Mountain Spotted Fever, for which doxycycline may be indicated. Doxycycline is preferred for its twice-a-day dosing low incidence of gastrointestinal side effects.
6. Although tetracyclines are not recommended during pregnancy, their use may be indicated for life-threatening illness. Adverse affects on developing teeth and bones are dose related, therefore, doxycycline might be used for a short course of therapy (7-14 days) prior to the 6<sup>th</sup> month of gestation. Please consult physician after the 6<sup>th</sup> month of gestation for recommendations.

## Tularemia (pneumonic, pleuropulmonary, typhoidal) Treatment Protocol

	Initial Therapy	Duration
<b>Adults</b>	Gentamicin 3-5 mg/kg/day (or in 3 divided doses, Q8 hrs) (IV or IM) <sup>1,2</sup> <b>OR</b>	<b>7 – 14 days</b> (If switching to oral doxycycline, duration of therapy should continue for a total of <b>21 days</b> )
	Doxycycline 100 mg IV BID <b>OR</b>	Estimated for <b>21 days</b> (May switch to oral doxycycline when clinically appropriate) <sup>3</sup>
	Ciprofloxacin 400 mg IV BID	Estimated for <b>10-14 days</b> (May switch to oral therapy when clinically appropriate)
<b>Children</b>	Gentamicin 6.0-7.5 mg/kg/day (or in 3 divided doses) (IV or IM) <sup>1,2</sup> <b>OR</b>	<b>7 – 14 days</b> (If switching to oral doxycycline, duration of therapy should continue for a total of <b>21 days</b> )
	Doxycycline <sup>4</sup> : <b>&gt; 8 yrs and &gt; 45 kg:</b> 100 mg IV BID <b>&gt; 8 yrs and ≤ 45 kg:</b> 2.2 mg/kg/day in 2 divided doses <b>≤ 8 yrs:</b> (same as <b>&gt; 8 yrs and ≤ 45 kg</b> ) <b>OR</b>	Estimated for <b>21 days</b> (May switch to oral doxycycline when clinically appropriate) <sup>3</sup>
	Ciprofloxacin <sup>5</sup> 15 mg/kg Q12hrs	Estimated for <b>10-14 days</b> (May switch to oral therapy when clinically appropriate)
<b>Pregnancy</b> <sup>6,7</sup>	Same as for non-pregnant adults	
<b>Immuno-compromised</b>	Same as for non-immunosuppressed adults and children	

1. Treatment of choice for tularemia is streptomycin. Streptomycin can be difficult to obtain; therefore gentamicin is often used and appears to be equally effective. Doxycycline is approved for the treatment of tularemia and is 90-100% absorbed after oral administration. This complete absorption may allow for its use in patients who can tolerate oral administration to complete the duration of therapy that is designated above.
2. The frequency of administration is left up to the discretion of the clinician, however, it should be noted that once-daily dosing of aminoglycosides is investigational. The manufacturers usually recommend that the daily dose be given in equally divided doses at 8-hour intervals; however, current evidence suggests that once-daily (single-daily) dosing of aminoglycosides is at least as effective as, and may be less toxic than, conventional dosing regimens employing multiple daily doses of the drugs.
3. To avoid relapses with shorter courses of therapy, longer courses of therapy are required when using oral doxycycline in *F. tularensis* due to doxycycline's bacteriostatic mechanism of action.
4. In 1991, the American Academy of Pediatrics amended their recommendation to allow treatment of young children with tetracyclines for serious infections, such as, Rocky Mountain Spotted Fever, for which doxycycline may be indicated. Doxycycline is preferred for its twice-a-day dosing low incidence of gastrointestinal side effects.
7. Ciprofloxacin dose should not exceed 1gram/day in children.
8. Aminoglycosides can cause fetal toxicity when administered to pregnant women, but potential benefits from use of the drug may be acceptable in certain conditions despite the possible risks to the fetus.
9. Although tetracyclines are not recommended during pregnancy, its use may be indicated for life-threatening illness. Adverse affects on developing teeth and bones are dose related, therefore, doxycycline might be used for a short course of therapy (7-14 days) prior to the 6<sup>th</sup> month of gestation. Please consult physician after the 6<sup>th</sup> month of gestation for recommendations.

## Plague Treatment Protocol

	Initial Therapy	Duration
<b>Adults</b>	Gentamicin 3-5 mg/kg/day (or in 3 divided doses, Q8 hrs) (IV or IM) <sup>1,2</sup> <b>OR</b>	<b>Estimated for 10 Days</b> (Switch to oral Doxy when clinically appropriate to complete <u>10 day</u> therapy)
	<b>Intravenous</b> Doxycycline 100 mg BID <b>OR</b>	
<b>Oral</b>	Doxycycline 100 mg BID	<b>Estimated for 10 Days</b>
<b>Children<sup>4</sup></b>	Gentamicin 6-7.5 mg/kg/day or in 3 divided doses (Q 8 hrs) <sup>2</sup> <b>OR</b>	<b>Estimated for 10 Days</b> (Switch to oral Doxy when clinically appropriate to complete 10 day therapy)
	<b>Intravenous</b> Doxycycline <sup>3</sup> : <u>&gt; 8 yrs and &gt; 45 kg</u> : 100 mg BID <u>&gt; 8 yrs and ≤ 45 kg</u> : 2.2 mg/kg/day in 2 divided doses <u>≤ 8 yrs</u> : (same as <u>&gt; 8 yrs and ≤ 45 kg</u> ) <b>OR</b>	
<b>Oral</b>	Doxycycline <sup>3</sup> : <u>&gt; 8 yrs and &gt; 45 kg</u> : 100 mg BID <u>&gt; 8 yrs and ≤ 45 kg</u> : 2.2 mg/kg BID <u>≤ 8 yrs</u> : 2.2 mg/kg BID	<b>Estimated for 10 Days</b>
<b>Pregnancy<sup>4,5</sup></b>	Same as for non-pregnant adults	
<b>Immuno-compromised</b>	Same as for non-immunocompromised adults and children	

1. Treatment of choice for plague is streptomycin. Streptomycin can be difficult to obtain; therefore gentamicin is often used as the drug of choice and appears to be effective.
2. The frequency of administration is left up to the discretion of the clinician, however, it should be noted that once-daily dosing of aminoglycosides is investigational. The manufacturers usually recommend that the daily dose be given in equally divided doses at 8-hour intervals; however, current evidence suggests that once-daily (single-daily) dosing of aminoglycosides is at least as effective as, and may be less toxic than, conventional dosing regimens employing multiple daily doses of the drugs.
3. In 1991, the American Academy of Pediatrics amended their recommendation to allow treatment of young children with tetracyclines for serious infections, such as, Rocky Mountain Spotted Fever, for which doxycycline may be indicated. Doxycycline is preferred for its twice-a-day dosing low incidence of gastrointestinal side effects.
4. Aminoglycosides can cause fetal toxicity when administered to pregnant women, but potential benefits from use of the drug may be acceptable in certain conditions despite the possible risks to the fetus.
5. Although tetracyclines are not recommended during pregnancy, its use may be indicated for life-threatening illness. Adverse affects on developing teeth and bones are dose related, therefore, doxycycline might be used for a short course of therapy (7-14 days) prior to the 6<sup>th</sup> month of gestation. Please consult physician after the 6<sup>th</sup> month of gestation for recommendations.

## References:

1. Bacterial Zoonoses Branch, DVVID, NCID, CDC: (December 14, 1999)
2. Emerging Bacterial and Mycotic Disease Branch, DBMD, NCID, CDC: (February 12, 1999)
3. AHFS Drug Information® 1999
4. Drugs Facts and Comparisons® 2000
5. 1997 Red Book, Report of Committee on Infectious Diseases, 24<sup>th</sup> Edition, American Academy of Pediatrics
6. Mandell, Douglas, and Bennett's, Principles and Practices of Infectious Diseases, 5<sup>th</sup> Edition
7. Control of Communicable Diseases Manual, 16<sup>th</sup> Edition, 1995
8. Inglesby TV, Henderson DA, Bartlett JG, et al. Anthrax as a Biological weapon, Medical and Public Health Management. JAMA 1999; 281(18): 1735-1745.
9. Red Book 2000, American Academy of Pediatrics, 25<sup>th</sup> Edition.
10. Blaser J, Konig C. Once-Daily Dosing of Aminoglycosides. Eur J Microbiol Infect Dis., 1995, 14:1029-1038.
11. Abramson J, Jon S, Givner LB. Rocky Mountain Spotted Fever. [Review] Ped Inf Dis J. 18(6):539-540, June 1999.
12. Inglesby TV, Dennis DT, Henderson DA, et al. Plague as a Biological Weapon, Medical and Public Health Management. JAMA 2000;283(17):2281-2290.
13. Use of Anthrax Vaccine in the United States: Recommendations of the Immunization Practices Advisory Committee (ACIP), MMWR, December 15 2000/Vol. 49/No. RR-15.
14. Inglesby TV, Dennis DT, Henderson DA, et al. Tularemia as a Biological Weapon, Medical and Public Health Management. JAMA 2001, In Press.